

Impact of Traditional Process on Hygienic Quality of *Soumbala* a Fermented Cooked Condiment in Burkina Faso.

Marius Kounbèsiounè Somda^{1*}, Aly Savadogo², François Tapsoba¹, Cheikna Zongo³, Nicolas Ouédraogo¹, Alfred Sabadénédyo Traoré^{1,2,3}

¹Laboratory of Microbiology and Biotechnology, Department of Biochemistry and Microbiology University of Ouagadougou.

²Laboratory of Food Technology, Department of Biochemistry and Microbiology University of Ouagadougou.

³Laboratory of Clinical Biochemistry and pharmacology, Department of Biochemistry and Microbiology University of Ouagadougou. 03 P.O Box: 7131 Ouagadougou 03, Burkina Faso.

Abstract

Soumbala is a African fermented locust beans used as a food condiment in Burkina that its quality could be affected due to hygienic practices. The present study aimed to show the implication of traditional process on hygienic quality of *soumbala*. *Soumbala* a fermented condiment was collected in six ethnical groups of Burkina Faso. A total of 120 samples were analysed using standard methods of microbiology. The microbial analysis showed a significant difference ($p < 0.05$) among microorganism groups following to the synthesized diagrams (A and B). Significant difference ($p < 0.05$) was established among microorganisms targeted in diagram A versus diagram B comparing separately the amount of *Bacillus sp.* ($18.77 \pm 0.29\%$ # $37.22 \pm 0.076\%$), *Micrococcus sp.* ($17.52 \pm 0.65\%$ # $10.68 \pm 0.38\%$) and *Staphylococcus sp.* ($19.63 \pm 0.35\%$ # $14.49 \pm 0.2\%$), Yeasts and Molds ($34.74 \pm 0.098\%$ # $37.6 \pm 0.26\%$), Total coliform ($9.1 \pm 0.16\%$ # 0). The results indicated that samples of *soumbala* contain various microorganisms due to the impact of processing.

Key words: Seeds, Process, Fermentation, *Soumbala*, Safety

Introduction

The need for additional protein supplies to promote sustainable livestock in most tropical areas has increased the search for indigenous wild legumes or condiments based seeds for least cost formulation and production. Two such legumes or condiments are *soumbala* and *bikalga*, commonly known as African locust bean, of tropical trees (*Parkia biglobosa* and *Hibiscus sabdariffa*) which are native to Africa and widely distributed in the savanna region (Adewusi et al. 1992). *Bikalga* and *Soumbala* are condiments of traditional uncontrolled alkaline fermentation of *Hibiscus sabdariffa* and *Parkia biglobosa* seeds respectively.

These food condiments are used and produced as major condiments in many African countries including Burkina Faso, Mali, Niger, Nigeria, Cameroon and Sudan among others. *Bikalga* is also called *Dawadawa botso* in Niger, *Datou* in Mali, *Furundu* in Sudan, *Mbuja* in Cameroon (Odufa 1985; Yagoub et al. 2004). *Bikalga* and *Soumbala* are produced by women and create an economical source (Ouoba et al. 2003). The seeds serve as a source of useful ingredients for consumption as *Iru* in Yoruba, have some anti-nutritional factors (ANFs) such as tannins, oxalate and hydrogen cyanide. ANFs were eliminated or reduced by application of heat and sprouting (Alabi et al. 2005). So fermentation has been reported to destroy some natural toxins which may occur in beans, improve the nutritive value, increase digestibility and enhance growth (Alabi et al. 2005; Bridget et al. 2004). The strains of *Bacillus* and *Staphylococcus* are implied fluently in the *soumbala* fermentation. *Bacillus* genus is involved in the initial fermentation of seeds characterized by the excretion of exopolysaccharides and lipopeptides (Savadogo et al. 2011).

* Corresponding author, mailing address:
50 33 73 73, Fax: (00226) 50 33 73 73,
E-mail: somdamarius@yahoo.fr

, Tel: (00226)

The *Staphylococcus* strains are responsible to the genesis of specific aromas. The excretion of lipopeptides can explain the use of these condiments as disinfectant in traditional medicine.

The *Bacillus subtilis* and *Bacillus pumilus* are implied in the deterioration of the indigestibles polysaccharides and oligosaccharides (arabinogalactan, stachyose) content in *Parkia biglobosa* seeds (Ouoba et al. 2003; Ouoba et al. 2007). Kim and Park (2009) showed that *Bacillus cereus* isolated in a fermented cooked that showed have a high anticancer effects against AGS Human Gastric Adenocarcinoma cells. The final product coming from the fermentation of seeds, can present itself in several aspects according to the preferences: flour, curds or wads. This product is used in the human food as condiment in sauces and can replace industrial soups.

The *soumbala* is condiment very consumed by some ethnic groups of Burkina Faso. An investigation permitted to target ethnic groups where its consumption is current. It is about the *Bissa*, *Dagara*, *Dafing*, *Mossi* and *Samos*. They are also consumed by *Peulh*, *Bwaba*, *bobo*, *Sambla*, *Sénoufo* (Ouoba et al. 2003).

This condiment is used preferentially in sauces and also in the preparation of the sauces of rice, *couscous* and other dishes basis of cereals. Several literature have been developed about fermented seed and related the extent of their physico-chemical properties and microbial aspect (Mohamadou et al. 2007; Ouoba et al. 2007; Parkouda et al. 2008; Oguntoyinbo et al. 2010; Parkouda et al. 2011).

The major problem of the set of the traditional process is located to the fluctuating quality of the different foods obtained. The processing and quality of *soumbala* is variable according *Mossi*, *Fulani*, *Bobo*, *Dagara*, *Samo*, *Bissa* groups. Indeed, the fermentation process which takes place spontaneously through the microflora development, can conduct to an organoleptic, microbiological or toxicological undesirable products (Rainbaul 1995).

This study was designed to value the impact of technological process of six ethnical groups on the microbial diversity of *soumbala*. It will be able to prevent pathogens contamination and improve the hygienic and safety quality.

Materials and Methods

The experiment was conducted in the Laboratory of Microbiology and Biotechnology at CRSBAN (University of Ouagadougou/Burkina Faso). The six processing kind of *soumbala* has been followed in the sites of production among ethnical groups as *Mossi*, *Fulani*, *Bobo*, *Dagara*, *Samo*, *Bissa* in Burkina Faso.

Sample collection Different samples of fermented condiments (*soumbala*) were taken to check the process effect on the microbial quality of these foods. Six (06)

process of *soumbala* production were studied. 120 samples of *soumbala* were collected on six (06) sites of production with seller women according to the ethnical groups mentioned in Table 1. In each site, 20 samples of *soumbala* were collected and maintained at + 4°C in isothermal box then sent to the laboratory for analysis.

Microbiological analysis The levels of total mesophilic flora, total coliform, *Escherichia coli*, *Staphylococcus sp*, *Micrococcus*, *Salmonella*, Yeast and Molds were determined by standards microbiology system using the methods of Zinnah et al (2007) and De Souza et al (2011). The prevalence rates and levels of presumptive were determined in 120 samples.

Preparation of sample. Each sample of *soumbala* was properly mixed to ensure homogenisation of the microorganisms present in the fermented product. 10 g of each of *soumbala* was introduced aseptically into 90 ml (1:10 dilution) of buffered Peptone water (0.85% w/v. Biomerieux, France) into stomacher blister and stored at ambient temperature for 30 min. The mixture introduce in a McCartney bottle was mixed with a vortex (Gemmy Industrial Corporation, Italy) for 5 min. A serial dilution of 10⁻¹ to 10⁻⁶ was made and 1 ml of each of dilution was pipetted in duplicate into appropriately marked Petri dishes.

Microbial numeration. All media used in this study were purchased from Biomerieux (France). The total number of microorganisms expressed as Colony Forming Unit (cfu) per gram of sample was determined by standard plate count.

- The total aerobic mesophilic flora (FAMT) was counted in Plate Count Agar (PCA), after 24 hours of incubation at 30°C under aerobic conditions.

- Eosin Methylen Blue medium (EMB) was carried out, to enumerate total coliforms and *E. coli*, after 48 hours of incubation, at 37°C and 44°C ±0.5°C respectively, under aerobic conditions.

- Yeasts and molds were counted in a Sabouraud after 3-5 days of incubation at 30°C, under aerobic conditions.

-*Salmonella* and *Shigella sp* was researched in a SS medium after 3-5 days of incubation at 37°C, under aerobic conditions.

- Suspected pathogenic *Staphylococcus sp*, were counted in the Chapman positive coagulase after 24 hours of incubation at 37°C, under aerobic conditions.

- *Micrococcus sp*, were counted in the Chapman medium after 24 hours of incubation at 37°C, under aerobic conditions.

- *Bacillus sp*, were counted in the Man Rogosa and Sharpe medium after 24 hours of incubation at 37°C, under aerobic conditions.

- Sulphite Reducing Bacteria (SRB) were screened with tryptone-sulfite neomycin broth after 20 hours incubation at 44°C±0.5°C.

Tabel 1. Diagrams of production of *soumbala*.

Process according to ethnic	Process Steps	Role of steps	Locality
Process of <i>mossi</i>	a , b, c, d, f, g, i, j	To obtain solid fermented product with pungent smell	Ouahigouya about 110 km of Ouagadougou
Process of <i>fulani</i>	a , b, c, d, f, g, i, j	To obtain solid fermented product with pungent smell	Dori about 400 km of Ouagadougou
Process of <i>samo</i>	a , b, c, d, e, f, g, h, i, j	To obtain soft to fell condiment with least pungent smell	Tougan about 300 km of Ouagadougou
Process of <i>bobo</i>	a , b, c, d, e, f, g, h, i, j	To obtain soft to fell condiment with least pungent smell	Bobo-Dioulasso about 365 km of Ouagadougou
Process of <i>dagara</i>	a , b, c, d, e, f, g, h, i, j	To obtain soft to fell condiment with least pungent smell	Dagara about 400 km of Ouagadougou
Process of <i>bissa</i>	a , b, c, d, f, g, i, j	To obtain solid fermented product with pungent smell	Garango about 200 km of Ouagadougou

Statistical analysis. Data obtained were expressed as means \pm standard deviation (SD). Statistical analysis was done using Stata/IC 10.0 from StataCrop LP. All enumeration values analyzed were transformed to log₁₀ values due to the abnormal nature of the data. A one way analysis of variance (ANOVA) was performed on enumeration values (duplicate means) obtained for each sample per medium. Comparison between mean values for the parameters considered was performed using the least significant difference test at 0.05 significance level.

Diagrams of production of *soumbala* Production of *soumbala* is a traditional process and involves many steps, including cleaning, boiling in alkaline conditions, pounding, fermentation steaming, drying and roasting. The different steps of process used for *soumbala* production have been regrouped in two diagrams.

The Figure 1 showed the similar steps process used by ethnical groups as *Mossi*, *Bissa*, *Fulani* while the Figure 2 denoted same steps in *Dagara*, *Bobo*, *Samo* ethnical groups. The two diagrams A and B showed that the main difference number of total mesophilic microorganisms coming from diagram A were most represented (log₁₀ (UFC/g): 9.42 \pm 0.03) was highest than diagram B (log₁₀ (UFC/g): 8.50 \pm 0.02) with a significant degree $p < 0.05$. Yeast and molds were count were significantly different (log₁₀ (UFC/g): 7.86 \pm 0.02 in diagram A versus log₁₀ (UFC/g): 6.63 \pm 0.06) in diagram B) with $p < 0.05$). A significant difference ($p < 0.05$) is noted concerning the presence of *Bacillus sp*, the diagram A (log₁₀ (UFC/g): 4.38 \pm 0.12) contain a less count than diagram B (log₁₀ (UFC/g): 6.46 \pm 0.02). And yet the notable remark was the presence of total coliform (log₁₀ (UFC/g): 2.13 \pm 0.06) in the *soumbala* provide from diagram A. The count of *Staphylococcus sp*

of process between ethnical groups reside in the method fermentation.

The soft to fell *soumbala* obtained in the diagram B was appreciated than one of diagram B and can be due to twice fermentation. During the fermentation, proteolysis followed by the degradation of carbohydrates seems to be the main biochemical and smell of seeds. This hypothesis is supported by the found of Ouoba, Parkouda, Diawara, Scotti and Varnam (2007).

They showed that proteolytic, lipolytic and amyolytic activities during the fermentation involve the change of fermented product.

Numeration of isolates microorganisms. The microbial numeration on different samples of *soumbala* was studied and is presented in Figure 3.

The results in Figure 3 showed that neither *Salmonella* nor Sulphite Reducing Bacteria were found in the analyzed *soumbala*. Total confirm were detected in Diagram A and empty in Diagram B. The samples were much contaminated by total aerobic microorganisms. It was observed that the and *Micrococcus sp* is higher from diagram A (log₁₀ (UFC/g): 4.58 \pm 0.15 and 4.09 \pm 0.02) than diagram B (log₁₀ (UFC/g): 2.51 \pm 0.05 at 1.85 \pm 0.11) with $p < 0.05$.

In accordance with the AFNOR limits (Association Française de Normalisation [AFNOR], 2009), all of samples investigate were acceptable in hygienic sight according to all enumerated microorganisms.




Figure 1. Diagram A relating to different steps of *soubala* production in Mossi, Bissa, Fulani





Figure 2. Diagram B relating to different steps of *soubala* production in Dagara, Bobo, Samo ethnic groups



Letters are used in alphabetical order to indicate variation value of microorganisms. The values are means \pm standard deviation of three replicates. Means with different alphabets on the same column group indicate are significantly different at $P < 0.05$ at 95% confidence level.

Figure 3. Number of isolates microorganisms following two diagrams of condiments production.

Microbial assessment. The distribution microbial following the diagram production of soumbala is presented in the Table 2. Results in Table 2 showed a significant difference ($p < 0.05$) between microbial communities presence in soumbala resulting of diagrams A and B of production. Significant difference ($p < 0.05$) was established

among microorganisms targeted in diagram A versus diagram B comparing separately the amount of *Bacillus* sp. ($18.77 \pm 0.29\%$ # $37.22 \pm 0.076\%$), *Micrococcus* sp. ($17.52 \pm 0.65\%$ # $10.68 \pm 0.38\%$) and *Staphylococcus* sp. ($19.63 \pm 0.35\%$ # $14.49 \pm 0.2\%$), Yeasts and Molds ($34.74 \pm 0.098\%$ # $37.6 \pm 0.26\%$), Total coliform ($9.1 \pm 0.16\%$ # 0).

Table 2. Percentage distribution of various microbial genera in *soumbala*

Microorganisms	Percentage of incidence (%)	
	Diagram A	Diagram B
Yeast and Mold	34.74 ± 0.098^a	37.6 ± 0.26^a
<i>Bacillus</i> sp	18.77 ± 0.29^{bc}	37.22 ± 0.076^a
Total coliform	9.1 ± 0.16^c	0
<i>Staphylococcus</i> sp	19.63 ± 0.35^b	14.49 ± 0.2^b
<i>Micrococcus</i> sp	17.52 ± 0.065^c	10.68 ± 0.38^c

Discussion

The *soumbala* collected on the sites of production in hot season are varied and undergo inopportune manipulations including the process of fermentation. These practices can reduce their quality. The *soumbala* is kept to an elevated ambient temperature. Temperature coupled an absence of cold chain causes a strong contamination (Gonfa et al. 2001). In the present work the microbiological quality of the *soumbala* examined is relatively acceptable.

Any sample was kept to limit to the quality of the international norms. It means unrespect of hygiene condition. The samples most contaminated are especially those of the diagram A due probably to the quality of water used during the production.

The detection of *S. aureus* and Sulphite Reducing Bacteria (SRB) is for their toxicological properties. So *S. aureus* is the leading species of the genus *Staphylococcus* implicated in food poisoning infection while SRB are a group anaerobic sporulating bacteria including *Clostridium*,

their spores resist after heat treatment and may cause damages after contaminated product consumption (De Souza et al. 2011). The contamination of samples pathogen may occur in *soumbala* processing or during the handling water and implement used also during the storage.

Similar shapes of contamination have been described on fermented products in several countries of Africa by some authors as Godefay and Molla (2000); Gonfa, Foster and Holzapfel (2001).

A number superior to 4.0 log (ufc/g) of *Mesophilic total* indicates deficiency condition of hygiene in the production or storage about Yamani, Al Kurdi, Haddadin and Robinson (1999). The microbiological quality of the condiment remained important for its conservation (Guinot-Thomas et al. 1995). The strong rate of yeasts and molds in *soumbala* samples could be explained as well as by the state of containers cleanliness used and the methods of transformation (Gadaga et al. 2000; Beukes et al. 2001).

The massive presence of yeasts and molds is also the expression of a strong outside contamination and a bad hygiene of the implement. The importance of *Staphylococcus aureus* and Total coliform in the fermented product could result to a lack of hygiene or a strong contamination of the raw material allowing germs to escape the bacteriocin produced by the *Bacillus* kind. *Staphylococcus aureus* and Total coliform are responsible to the food damage. The proliferation of the microorganisms on the *soumbala* could be explained by the necessary mineral and nitrogenous elements found in abundance on the raw material and used by these microorganisms (Somda et al. 2011).

Conclusion

The results showed that *soumbala* contain various microorganisms. The presence of the indicator germs of hygiene lack shows that the *soumbala* consumed by an important part of the population can especially contribute to reduce the quality and time conservation then create economical impact.

Acknowledgement

We need to thank Pr Alfred S. Traoré responsible of Research Center in Biological Food and Nutrition Sciences (CRSBAN) and also producer women of *soumbala* for their collaboration.

References

Adewusi SRA, Orisadare BA, Oke OL. 1992. Studies on weaning diets in Nigeria. Two protein sources. *Plant Food Human Nutr.* 42: 183-192.

Alabi DA, Akinsulire OR, Sanyaolu MA. 2005. Qualitative determination of chemical and nutritional composition of *Parkia biglobosa*. *Afric. J. Biotechnol.* 4: 812-815.

Association Française de Normalisation (AFNOR). 2009. Normes Agroalimentaires. <http://www.afnor.org/secteurs/activite/agroalimentaire/normes/liste-desnormes>.(accessed January 15, 2014).

Beukes EM, Bester B H, Mostert J F. 2001. The microbiology of South African traditional fermented milks. *Internatio. J. of Food Microbiol.* 63: 189-197.

Bridget OO, Olumuyiwa SF, Bolanie OA, Adewusi SAA. 2004. Biochemical changes in Africa locust beans (*Parkia biglobosa*) and melon (*Citrullus vulgaris*) seeds during fermentation to condiment. *Pakistan J. Nutrit.* 3: 140-145.

De Souza C, Ameyapoh Y, Karou SD, Anani KT, Kpodar ML, Gbeassor M. 2011. Assessing Market-Sold Remedies in Lomé (Togo) for Hygienic Quality. *Biotechnol. Research Internatio.* pp:1-5.

Gadaga TH, Mutukumira AN, Narvhus JA. 2000. Enumeration and identification of yeasts from Zimbabwean traditional fermented milk. *Internatio. Dairy J.* 10: 459- 466.

Godefay B, Molla B. 2000. Bacteriological quality of raw cow's milk from four dairy farms and a milk collection centre in and around Addis Ababa. *Berlin. Munchener Tierar. Wochen.* 113: 276-278.

Gonfa A, Foster HA, Holzapfel WH. 2001. Field survey and literature review on traditional fermented milksabdariffa seeds in Cameroon. *J. of Food Technol.* 5: 164-168.

Odufa SA. 1985. Biochemical changes during "iru"fermentation. *J. of Food Technol.* 20: 295-303.

Oguntoyinbo FA, Huch M, Cho GS, Schillinger U, Holzapfel WH, Sanni AI, Franz CM. 2010. Diversity of *Bacillus* species isolated from okpehe, a traditional fermented soup condiment from Nigeria. *J. of Food Protect.* 73: 870-878

Ouoba LII, Parkouda C, Diawara B, Scotti C, Varnam AH. 2007. Identification of *Bacillus* spp from Bikalga, fermented seeds of *Hibiscus sabdariffa*: phenotypic and genotypic characterization. *J. of Appl. Microbiol.* 104: 122-131.

Ouoba LII, Cantor MD, Diawara B, Traore AS, Jakobsen M. 2003. Degradation of African locust bean oil by

- Bacillus subtilis and Bacillus pumilus isolated from soumbala, a fermented African locust bean condiment. *J. of Appl. Microbiol.* 95 (4): 868-873.
- Parkouda C, Diawara B, Lowor S, Diako C, Kwesi-Saalia F, Annan NT, Jensen JS, Tano-Debrah K, Jakobsen M. 2011. Volatile compounds of maari, a fermented product from baobab (*Adansonia digitata* L.) seeds. *Afric. of J. Biotechnol.* 10 (20): 4197-4206.
- Parkouda C, Diawara B, Ouoba LII. 2008. Technology and physico-chemical characteristics of Bikalga, alkaline fermented seeds of *Hibiscus sabdariffa*. *Afric. of J. Biotechnol.* 7: 916-922.
- Rainbault A. 1995. Bactéries lactiques et fermentation du manioc. In: Transformation alimentaire du manioc. Eds T. Agbor Egbe, A. Brauman, D. Griffon and S. Trèche. Paris, ORSTOM. pp. 258-275.
- Savadogo A, Tapi A, Chollet M, Wathelet B, Traoré AS, Jacques P.H. 2011. Identification of surfactin producing strains in Soumbala and Bikalga fermented condiments using Polymerase Chain Reaction and Matrix Assisted Laser Desorption/Ionization-Mass Spectrometry methods. *Internatio. J. Food of Microbiol.* 151:299-306.
- Seo HR, Kim JY, Kim JH, Park KY. 2009. Identification of *Bacillus cereus* in a Chungkukjang that Showed High Anticancer Effects Against AGS Human Gastric Adenocarcinoma Cells. *J. of Medici. Food.* 31: 1274-1280.
- Somda KM, Savadogo A, Barro N, Thonart P, Traoré AS. 2011. Effect of mineral salts in fermentation process using mango residues as carbon source for bioethanol production. *Asi. of J. Industri. Engineer.* 3(1): 29-38.
- Yagoub AEGA, Mohamed BE, Ahmed AHR, El-Tinay AH. 2004. Study on Furundu, a traditional sudanese fermented roselle (*Hibiscus sabdariffa*) seed: effect on in vitro protein digestibility, chemical composition and functional properties of the total proteins. *J. of Agricult. Food Chemist.* 52: 6143-6150.
- Yamani MI, Al-Kurdi LMA, Haddadin MSY, Robinson RK. 1999. A simple test for the detection of antibiotics and other chemical residues in ex-farm milk. *Food Control.* 10: 35-36.
- Zinnah MA, Bari MR, Islam MT, Hossain MT, Rahman MT, Haque MH, Babu SAM, Ruma RP, Islam MA. 2007. Characterization of *Escherichia coli* isolated from samples of different biological and environmental sources. *Banglad. J. of Veterin. Med.* 5 (2): 25-32.