

Prevalence of Salmonella Serovars Isolated from Turkey Carcasses and Giblets in Meknès-Morocco

El Allaoui 1*, F. Rhazi Filali 2, B. Bouchrif 3, B. Karraouan4 and Ameer N 5

1,2 Equipe Microbiologie et Santé, Département de Biologie, Laboratoire de Chimie Biologie Appliquées à l'Environnement, Université Moulay Ismail Faculté des Sciences, B.P.11201 Zitoune Meknès, Maroc.

3,4Institut Pasteur du Maroc, Laboratoire de microbiologie et de l'hygiène des aliments et des eaux

5 Département de Microbiologie eaux aliments, Institut National d'Hygiène Rabat

Abstract

The present study was conducted to determine the prevalence and the serotypes involved the virulence gene (InvA and SpvC) of Salmonella isolates recovered from the raw meat and giblets (liver and gizzard) of the turkey in various outlets in the Moroccan market. From November 2011 to November 2012 a total of 192 samples of turkey meat (included 48 breast, 48 legs, 48 gizzards and 48 livers) were collected every ten days from retail outlets in Meknès. Of these, 48 were from popular market, 48 from artisanal slaughterhouses, 48 from poulterers' shops and 48 from a supermarket at Meknes, Morocco. Of the total of 192 samples examined, 24.5% (47/192) were contaminated with Salmonella. Out of the total 48 samples analysed from popular market, 19 (40.42%) proved to be Salmonella positive whereas from 48 samples obtained from traditional slaughterhouses and 48 from poulterers' shops 14 (29.87%) and 8 (17%) contained Salmonella, respectively. Compared to other outlets, a low level of Salmonella contamination was found in samples obtained from Supermarket 6 (12.7%). Among the 47 Salmonella isolates, 6 different serotypes were identified of which S. Saintpaul (46.8%) was the most frequent, followed by S. Agona (17%) and S. Kentucky (17%), S. Typhimurium (8.5%), S. Infantis (6.3 %) and S. Bredeney (4.2 %). The high level of contamination, especially in popular market and artisanal slaughterhouses of turkey meat and giblets with Salmonella observed in this paper indicates the need for an improvement in the microbiological quality of retail turkey. Examination of Salmonella for invA gene was detected in all the strains (n =47), only three isolates were positive for the gene SpvC: S. Agona, S. Kentucky and S. Infantis.

Key words: Salmonella, Turkey, Retail outlets, Meknès, InvA, SpvC, Morocco

Introduction

Salmonella infections occur worldwide in both developed and developing countries and are a major contributor to morbidity and economic costs (Antoine et al. 2008). According to the World Health Organisation (WHO), there are about 17 million cases of acute gastroenteritis or diarrhea annually due to non-typhoidal salmonellosis with 3 millions deaths (Rabsch et al. 2001). In some industrialized countries like the United States and Great Britain country, turkey meat is responsible for twice as cases of salmonellosis in humans that products made with chicken (Bryan et al.1998).

In several countries, a high level of Salmonella contamination in chicken carcasses and giblets from processing plants or retail markets has been reported (Arumugaswamy et al.1995; Carraminana et al.1997; Dominguez et al. 2002). Previous works undertaken in Morocco indicated the presence and distribution of Salmonella in poultry farms (Chaiba, 2010), Turkey meat and meat products (Cohen et al.2007; karraouan et al.2010), selected food items (Bouchrif et al.2009) and man (Ammari et al. 2009). In Morocco, Salmonella, Staphylococcus Aureus, and Clostridium Perfringens are reported to cause 42.8, 37 and 1.7% of food poisoning, respectively (Department of Epidemiology, 2009). Although the declaration and recording of 12 % Salmonella cases remain underreported, Salmonella is the major cause of food poisoning in Morocco (Rouahi et al. 1998), In France is

* Corresponding author. mailing address: Department of biology, College of sciences, Moulay Ismail University BP 11 201. Zitoune. Meknès. Morocco, Tel: +212668479701, Email: alaouixsaraa@hotmail.com

responsible for collective food poisoning with approximately 65% (Haeghebaert et al. 2009) of cases and of 95% in the United States of America (Mead et al.1999) Four kinds of meat turkey outlets are used in Morocco: popular market, artisanal slaughterhouses, poulterers' shops and supermarket. They differ from each other by the level of hygiene, diet cold which are subject carcasses (ambient temperatures, refrigeration, freezing) .At popular market and artisanal slaughterhouses the conditions of slaughter and sale of the product are faulty (Amara et al. 1994) , This kind of poultry is often sold in parts and the selling can take time, during which the carcasses are displayed at ambient temperatures during the day and put in the refrigerator for the night (Amara et al. 1994; Aymar et al. 1998). On the contrary, poulterers'shops and supermarket ensure the slaughter, storage and sale of poultry meat under good hygienic conditions (Direction d'élevage, 2007).

Currently, there is limited information regarding the prevalence of Salmonella in poultry, especially turkey, in Morocco. Therefore, the present study was conducted to determine the Salmonella prevalence and the serotypes involved, the virulence gene (InvA and SpvC) of Salmonella isolates recovered from the raw meat of the turkey in various outlets in Meknes.

Materials and Methods

Samples. Between October 2011 and October 2012, a total of 192 samples of turkey meat (included 48breast, 48 legs, 48 gizzards and 48 livers) were collected every ten days from retail outlets in Meknès. Of these, 48 were from popular market, 48 from artisanal slaughterhouses, 48 from poulterers'shops and 48from a supermarket at Meknes, Morocco. Each sample (approximately 50 g) was placed in a separate sterile plastic bag. Samples were transported to the laboratory immediately after collection in an ice chest and microbiological analysis was carried out immediately.

Isolation and identification of *Salmonella*.

Bacteriological analysis is conducted according to the AFNOR standard (NF U 47-100). Suspected colonies for *salmonella* were inoculated in Hanja Kligler (Biokar Diagnostic, France) at 37°C for 18 to 24 hours, in Urea Indol (bio Mérieux^R SA, France) at 37°C for 2 to 4 hour and testing wit oxydas disc (In vitro Diagnostic, USA), in citrate (Oxoid, England), in mannitol (Biokar Diagnostic, France), in lysine (SCHARLAB, Barcelona), and wit an ONPG disc (Oxoid Limtedt, England) for biochemical testing and presumptive identification. All isolates were biochemically identified by using the API 20^E system (bio Mérieux^R SA, France) and then serotyped by slide agglutination test using Salmonella polyvalent O and H antisera (Diagnostic Pasteur, Paris,France).

Detection of virulence genes. The Salmonella isolates were analyzed by PCR to detect the presence of virulence genes *invA* and *spvC*. PCR amplification was performed

using primer pairs described by Bhatta et al (2007) [34] (F-5'tatcgccacgttcgggcaa3' and R-5'tcgaccgtcaaaggaacc3' and F-5'cggaatacatcaata3' and R-5'cccaaaccatactactctg3' respectively).

The amplicon sizes of *invA*, *spvC* were 275 bp and 669 bp respectively. Amplification was performed in a 25 µl final volume, with a reaction mixture containing 1 µl bacterial DNA; 5 µl green GO Taq buffer (5x); 100 µM each eoynucleoside triphosphates (dNTPs), 0.125 µM each primers, and 0.5 U GO Taq DNA polymerase (Bio-Rad). Amplification was conducted in the thermocycler (Verity ,Bio-Rad). The PCR cycling program of the virulence gene *invA/spvC* consisted of denaturation at 94 °C for 4 min, followed by 40 cycles of 94 °C for 30 s, 52°C for 30 s, 72 °C for 45 s, and a final extension period at 72 °C for 7 min. PCR products (4 µl) were resolved by electrophoresis in 1.5–2% (w/v) agarose gels and visualized under ultraviolet transillumination after ethidium bromide staining. A wide-range molecular-weight DNA marker (100-bp DNA ladder, Promega) was used on each gel as a standard. Salmonella Typhimurium ATCC 14028 was used as control for all PCR detection.

Results and Discussion

Of the total of 192 samples examined, 24.5% (47/192) were contaminated with *Salmonella* (Table). Out of the total 48 samples analysed from popular market, 19(40.42%) proved to be *Salmonella* positive whereas from 48 samples obtained from traditional slaughterhouses and 48 from poulterers'shops 14(29.87%) and 8 (17%) contained *Salmonella*, respectively. Compared to other outlets, a low level of *Salmonella* contamination was found in samples obtained from Supermarket 6 (12.7%). Among the 47 *Salmonella* isolates, 6 different serotypes were identified of which *S. Saintpaul* (46.8%) was the most frequent, followed by *S. Agona* (17%) and *S. Kentucky* (17%), *S. Typhimurium* (8.5%), *S. Infantis* (6.3 %) and *S. Bredeney* (4.2 %). As shown in Table 2, a high level of *Salmonella* contamination was found in turkeybreast (44.6%) and gizzard (23.4 %), followed by livers (17%) and legs (12.7%). Contamination of turkey carcasses (28.6%) was higher than those of rates of turkey giblets (20, 2 %).

Table 1. *Salmonella* isolated from retail outlets

Sample from	Number of samples		
	Examined	Positive	%
Popular Market	48	19	40,42
Artisanal slaughterhouses	48	14	29,87
Poulterers'shpos	48	8	17
Supermarket	48	6	12,7

Table 2. Distribution of *Salmonella* serotypes in turkey meat and giblets (n=192)

Serotype	Number of positive Samples				Total 192
	Breast n=48	Legs n=48	Liver n=48	Gizzard n=48	
<i>S. Saintpaul</i>	10	2	4	6	22
<i>S. Agona</i>	4	1	1	2	8
<i>S. Kentucky</i>	4	2	1	1	8
<i>S. Typhimurium</i>	1	1	1	1	4
<i>S. Bredeney</i>	-	-	1	1	2
<i>S. Infantis</i>	2	1	-	-	3
Total	21	6	8	11	47

Examination of *Salmonella* for *invA* gene was detected in all the strains (n =47), only three isolates were positive for the gene *SpvC*: *S. Agona*, *S. Kentucky* and *S. Infantis* (Figure 1).




Figure1. Agarose gel electrophoresis of amplicons generated by simple PCR using primers specific for *Salmonella* virulence genes. Line 1: 275-bp *invA* amplicon; Line 2: 669-bp *SpvC* amplicon, C⁺ and T⁺: Positive control (*S. Typhimurium* penta-resistant ACTeStSul kind) C⁻, T⁻: negative control, M: 100-bp DNA ladder.

Discussion

The contamination rate of turkey carcasses and giblets observed in this study (24.94%) was in agreement with those reported by Karraouan *et al.* (2010) (20.3% in raw turkey minced meat) and Chaiba and al. (2009) (20.83% in chicken carcasses, chicken parts and giblets) in Morocco, by Cook *et al.* (2009) in Ontario (24% in turkey pieces), by Arslan and Eyi (2010) in Turkey (29.3% , poultry from delicatessens). However, Beli *et al.* (2001) reported a low prevalence of *Salmonella* in turkey meat in Albania (8.2%), Jordan *et al.* (2006) in Ireland (3.1%), in USA 2, 6 %, and Zhao *et al.* (2001) in the UK (5.6%). nevertheless, Bentley

(1984) reported a higher prevalence of *Salmonella* in turkey meat in Canada (68.8%).

As shown in Table 1, the contamination rate in poulterers' shops carcasses (17%) was higher than in supermarket (12.7%) , possibly due to the greater use of "use by" dates by the supermarkets and also of packaging that would prevent further cross contamination between samples. The results coincide from those obtained by Plummer *et al.* (1995), who detected a lower number of salmonella contaminated carcasses from supermarkets (18.6%), than poulterers' shops (24.5%). This comparison should be made with caution because several factors must be taken into account when making such comparisons, including differences in country and origin, type of meat samples, sampling seasons, slaughterhouse sanitation, and isolation methods. The high prevalence rates reported here might be due to a combination of the low quality of turkey carcasses used, especially in popular market and artisanal slaughterhouses, indeed cross-contamination of *Salmonella* from giblets to carcass could occur during handling, processing, packing and distribution. The packing of giblets with the carcass observed in this study could have also contributed to increase *Salmonella* cross-contamination. In addition to these, scalding water can become contaminated with *Salmonella* from faeces, plucking equipment, cages and floors. Workers can spread the contamination during retailing (Arumugaswamy *et al.*1995; Uyttendaele *et al.*1998). Rupture of the intestine could also occur during evisceration and pooling giblets might lead to cross-contamination of carcasses and other turkey parts. The process of conventional defeathering has been showed to play an important role in contamination of a high number of turkey carcasses (Clouser *et al.* 1995). Two out of the sex serotypes detected in the present study (*S. Infantis* and *S. Typhimurium*) are in the top 5 most frequent serotypes associated with human salmonellosis in the European Union in the last years (EFSA, 2011).Whereas *S. Kentucky* was the most common serovar found in Ontario during a sampling period from 2005 to 2006 (CIPARS, 2007) . During the period 2002–2005, we reported 17 cases of salmonellosis in French travelers returning from northeast and eastern Africa, from whom *S. enterica* serotype

Kentucky isolates resistant to ciprofloxacin with were recovered (Weill *et al.* 2006). This serovar was detected in Morocco, between 2005 and 2008, in raw turkey minced meat to 20.5% (Karraouan *et al.* 2010). *S. Saintpaul*, the most frequently isolated (46.8 %) in this study, has been associated with foodborne outbreaks including one due to contaminated paprika (Guinee *et al.* 1961).

S. Saintpaul was detected in fattening turkeys in 12 countries, reflecting the wide spread of this serovar (Janine *et al.* 2010). According to Enter-Net reports (data on *Salmonella* human isolates identified by European national reference centers), for the last quarter of the year 2006 (Janine *et al.* 2010). *Salmonella enterica* serovar Agona was first identified in Ghana (Guinee *et al.* 1961). Since then, this serovar has been reported in many countries worldwide in both humans and animals (Clark *et al.* 1973).

Invasion gene operon, *invA* was detected in all *Salmonella spp.* isolates in our study. This gene is essential for full virulence in *Salmonella* and is thought to trigger the internalization required for invasion of deeper tissue (Khan *et al.* 1999). There are studies reporting the detection of this gene in all *Salmonella spp.* isolates (Zahraei *et al.* 2006; Nashwa *et al.* 2009; Karraouan *et al.* 2010). Another study on finding the *SpvC* gene in different serovars isolated from bovine *Salmonella* and pathological products humans by PCR revealed a frequency of 28% (21/75) and none of the isolates of *Salmonella* serotypes (Heidelberg ; n = 17), (Infantis ; n = 10), (Schwarzengrund ; n = 11) isolated from broilers were positive for the gene *SpvC* (Abouzeed *et al.* 2000).

Conclusion

The high level of contamination of turkey meat and giblets with *Salmonella* observed in this paper indicates the need for an improvement in the microbiological quality of retail turkey. There is also a need for a comprehensive epidemiological study and control of *Salmonella* contamination at various levels of turkey production and retail outlets in Morocco.

References

- Abouzeed YM, Hariharana H, Poppeb C and Kibengea FSB. Characterization of *Salmonella* isolates from beef cattle, broiler chickens and human sources on Prince Edward Island. *Comparative Immunology, Microb. Infect. Dis*, 23: 253-266. 2000.
- Amara A, Badoum M, Faid M and Bouzoubaa K. Microbial contamination of poultry slaughtered in traditional shops in Morocco. *Microbiol. Aliment Nutr*, 12 : 323-327. 1994.
- Aymar J. Appréciation de la qualité bactériologique des carcasses de la volaille préparées dans un abattoir avicole industriel à Rabat, Thèse de Doctorat Vétérinaire, Institut Agronomique et Vétérinaire Hassan II, Rabat .1998.
- Ammari S, Laglaoui A , En-nanei L, Bertrand S, Wildemauwe C, Barrijal S and Abid M . Isolation, drug resistance and molecular characterization of *Salmonella* isolates in northern Morocco. *J.Infect Developing Countries*, 3(1): 41-49. 2009.
- Anonymous. Fisa. Documentations & statistiques. 2008.
- Antoine ST, Annaelle K and Anne B. Epidemiological analysis of *Salmonella enterica* from beef sampled in the slaughterhouse and retailers in Dakar (Senegal) using pulsed-field gel electrophoresis and antibiotic susceptibility testing. *J Food Microbiol*, 123: 191-197. 2008.
- Antunes P, Reu C, Sousa JC, Peixe L and Pestana N. Incidence of *Salmonella* from poultry ucpodts and their susceptibility to antimicrobial agents. *Int. J. Food Microbiol*, 82: 97-103.2003.
- Arslan S and Eyi A. Occurrence and antimicrobial resistance profiles of *Salmonella* species in retail meat products. *Journal of Food Protection*, 73: 1613-1617. 2010.
- Arumugaswamy RK, Rusul G, Abdul Hamid SN and Cheah CT. Prevalence of *Salmonella* in raw and cooked foods in Malaysia. *Food Microbiol*, 12: 3-8.1995.
- Beli E, Telo A and Duraku E. *Salmonella* serotypes isolated from turkey meat in Albania. *Int. J. Food Microbiol*, 63: 165-167. 2001.
- Bhatta DR, Bangtrakulnonth A, Tishyadhigama P, Saroj SD, Bandekar JR and Hendriksen RS. Serotyping, PCR, phage-typing and antibiotic sensitivity testing of *Salmonella* serovars isolated from urban drinking water supply systems of Nepal. *Letters in Applied Microbiology*, 44:588-594. 2007.
- Bentley AH. The *Salmonella* situation in Canada. In: Proc. Int. Symp. on Salmonella, American Association Avian Pathologists., Snoeyenbos GH (Ed.) , 54-63. 1984.
- Bouchrif B, Paglietti B, Murgia M, Piana A, Cohen N, Ennaji MM, Rubino S and Timinouni M. Prevalence and antibiotic-resistance of *Salmonella* isolated from food in Morocco. *J Infect. Dev. Ctries*, 3(1):35-40. 2009.

- Bryan FL, Michanie SC, Alvarez P and Paniagua A. Critical control points of street-vended foods in Dominican Republic. *Journal of Food Protection*, 51: 373-383. 1988.
- Capita R, Alonso-Calleja C, Prieto M. Prevalence of *Salmonella* enterica serovars and genovars from chicken carcasses from slaughterhouses in Spain. *Journal of Applied Microbiology*, 103: 1366-1375. 2007.
- Carraminana JJ, Yanguela J, Blanco D, Rota C, Agustin AI, Arino A and Herrera A. *Salmonella* incidence and distribution of serotypes throughout processing in a spanish poultry slaughterhouse. *J. Food Prot*, 60: 1312-1317. 1997.
- Chaiba A. Impact des pratiques de production de poulet de chair à Meknès sur la qualité bactériologique, l'antibiorésistances et les résidus d'antibiotiques dans les produits aviaires finis, *Thèse de Doctorat National*, Université Moulay Ismail, Faculté des Sciences de Meknès, Maroc. 2011.
- CIPARS. Canadian Integrated Program for Antimicrobial Resistance Surveillance Annual Report. Public Health Agency of Canada, Guelph, Ontario, Canada. 2007.
- Weill FX, Bertrand S, Guesnier F, Baucheron S, Grimont PAD and Cloeckaert A. Ciprofloxacin-resistant *Salmonella* Kentucky in travelers. *Emerg Infect Dis*, 12:1611-2. 2006.
- Lehmacher A, Bockemuhl J and Aleksic S. Nationwide outbreak of human salmonellosis in Germany due to contaminated paprika and paprika-powdered potato chips. *Epidemiol Infect*, 115:501-511. 1995.
- Clouser CS, Doores S, Mast MG and Knabel SJ. The role of defeathering in the contamination of turkey skin by *Salmonella* species and *Listeria monocytogenes*. *Poult. Sci*, 74:723-731. 1995.
- Cohen N, Ennaji H, Bouchrif B, Hassar M and Karib H. Comparative Study of Microbiological Quality of Raw Poultry Meat at Various Seasons and for Different Slaughter Processes in Casablanca (Morocco). *Poultry Science Association, Inc. J. Appl. Poult. Res*, 16: 502-508. 2007.
- Cook A, Reid-Smith R, Irwin R, McEwen SA, Valdivieso-García A and Ribble C. Antimicrobial resistance in *Campylobacter*, *Salmonella*, and *Escherichia Coli* isolated from retail turkey meat from Southern Ontario, Canada. *Journal of Food Protection* 72: 473-481. 2009.
- Department of Epidemiology. Foodborne Disease Outbreak Reports, Searchable Data 2000-2005. Ministry of Public Health, Rabat, Morocco. 2005.
- Direction de l'élevage. Rabat, Maroc. Situation du secteur avicole à la veille de l'application de la loi 49/99. 2007.
- Dominguez C, Gomenz I and Zumalacarregui J. Prevalence of *Salmonella* and *Campylobacter* in retail chicken meat in Spain. *Int. J. Food Microbiol*, 72:165-168. 2002.
- Jerngklinchan J, Koowatananukul C, Daengprom K and Saitanu K. Occurrence of *salmonella* in raw broilers and their products in Thailand. *J. Food Prot*, 57:808-810. 1994.
- EFSA. The European Union summary report on trends and sources of zoonoses, zoonotic agents and foodborne outbreaks in 2009.
- EFSA Journal 9 (3), 2090-2477 Available at: <http://www.efsa.europa.eu/en/efsajournal/doc/2090.pdf> Date last accessed: May 4, 2011.
- Guinee P, Kampelmacher E and Willems H. Six new *Salmonella* types, isolated in Ghana (*S. volta*, *S. agona*, *S. wa*, *S. techimani*, *S. mampong* and *S. tafo*). *Antonie Van Leeuwenhoek*, 27:469-72. 1961.
- Haeghebaert S, Sulem P, Deroudille L, Vanneroy-Adenot E, Bagnis O, Bouvet P, Grimont F, Brisabois A, Le Querrec F, Hervy C, Espie E, de Valk H and Vailant V. Two outbreaks of *Salmonella* Enteritidis phage type 8 linked to the consumption of Canal cheese made with raw milk, France, 2001. *Eurosurveillance*, 8:151-156. 2001.
- ICMSF. International Commission on Microbiological Specifications for Foods. Poultry and poultry products. In: *Microorganisms in Foods. 6. Microbiological Ecology of Food Commodities*. Blackie Academic & Professionals, London. 1998.
- Janine B, Irene R, Andreas S, Annemarie Ka's, Reiner H and Beatriz G. A Predominant Multidrug-Resistant *Salmonella* enterica Serovar Saintpaul Clonal Line in German Turkey and Related Food Products. *applied and environmental microbiology*, 3657-3667. 2010.
- Jordan E, Egan J, Dullea C, Ward J, Mc Gillicuddy K, Murray G, Murphy A, Bradshaw B, Leonard N, Rafter P, Mc Dowell S. *Salmonella* surveillance in raw and cooked meat and meat products in the

- Republic of Ireland from 2002 to 2004. *Int. J. Food Microbiol*, 112: 66-70. 2006.
- Karraouan B, Fassouane A, EL Ossmani H, Cohen N, Charafeddine O et Bouchrif B. Prévalence et gènes de virulence des *Salmonella* isolées des viandes hachées crues de dinde à Casablanca (Maroc). *Revue Méd. Vét*, 161, 3: 127-132. 2010.
- Khan AA, Nawaz MS, Khan S and Sernigelia CE. Detection of Multidrug Resistant *Salmonella* Typhimurium DT104 by Multiplex polymerase chain reaction. *FEEMS Microbiol. Lett*, 182: 355-360. 1999.
- Mead PS, Slutsker L, Diets V, McCaig LF, Bresee JS, Shapiro C, Griffin PM and Tauxe RV. Food related illness and death in United States *Emerg Infect Dis*, 5:607-625. 1999.
- Nashwa HM, Mahmoud AH, Sami and Adawy S. Application of Multiplex Polymerase Chain Reaction (M-PCR) for Identification and Characterization *Salmonella* enteritidis and *Salmonella* typhimurium, 5(12): 2343-2348. 2009.
- Plummer RAS, Blissett ST and Dodd CER. *Salmonella* contamination of retail chicken products sold in the UK. *J. Food Prot*, 58:843-846. 1995.
- Rabsch W, Tschape, H and Baumler, A J. Non-typhoidal Salmonellosis: Emerging problems. *Microbes and Infection*, 3: 237-247. 2001.
- Rouahi N, Zouhdi M, Benabderrazek F, Boudhan A, Hamid K, Driss L, Zidouh A, Benkadour k . Mahjour J, Elyachoui M et Alaoui MA. Analyse des données des trois dernières années sur les salmonelloses au Maroc (1995-1997), *Biologie infection Tome IV*: 3-10. 1998.
- Yashoda KP, Sachindra NM , Sakhare PZ and Rao DN . Microbiological quality of broiler chicken carcasses processed hygienically in a small scale poultry processing unit. *J. Food Quality*, 24: 249-259. 2001.
- Zahraei T, Mahzoonae MR, and Ashrafi A. Amplification of *invA* gene of *Salmonella* by polymerase chain reaction (PCR) as a specific method for detection of *Salmonella*. *J. Fac. Vet. Med. Univ. Tehran*, 61(2): 195-199. 2009.
- Zhao C, Ge B, Villena JD, Sulder R , Yeh E, Zhao S, White DG, Wagner D and Meng J. Prevalence of *Campylobacter* spp. , *Escherichia Coli*, and *Salmonella* serovars in retail chicken , turkey, pork and beef from Grater Washington, D.C, Area. *Appl Environ Microbiol*, 67:5431-5436. 2001.