

Insecticidal Activity *Aegle marmelos* (L.) Correa Essential Oil Against Four Stored Grain Insect Pests

Rajesh Kumar¹, Ashok Kumar¹, Chandra Shekhar Prasad¹, Nawal Kishore Dubey^{1*} and Raju Samant²

¹ Centre of Advanced Study in Botany, Banaras Hindu University, Varanasi, India

² Department of Entomology, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi, India

Abstract: Experiments were carried out to determine the potential of using essential oil from leaves of *Aegle marmelos* to control insect infestation of stored gram from *Callosobruchus chinensis* (L.) (Bruchidae) and wheat from *Rhyzopertha dominica* (F.) (Bostrychidae), *Sitophilus oryzae* (L.) (Curculionidae) and *Tribolium castaneum* (Herbst) (Tenebrionidae). After introducing the test insects, stored gram and wheat samples were fumigated with essential oil of *Aegle marmelos* at 500 µg/mL (ppm). The oil significantly enhanced feeding deterrence in insects and reduced the grain damage as well as weight loss in fumigated gram and wheat samples infested with all insects except *T. castaneum*. The essential oil at different doses significantly reduced oviposition and adult emergence of *C. chinensis* in treated cowpea seeds. The oil protected stored gram from *C. chinensis* and wheat from *R. dominica* and *S. oryzae* for two years. Limonene (88 %) was found to be the major component in the oil through GC-MS analysis. Regression analysis of data on individuals in treated cowpea confirmed that significant reduction of oviposition and adult emergence of *C. chinensis* decreased with increase in doses. The findings emphasize the efficacy of *A. marmelos* oil as fumigant against insect infestations of stored grains and strengthen the possibility of using it as an alternative to synthetic chemicals for preserving stored grains.

Key words: *Aegle marmelos*, *Callosobruchus chinensis*, *Rhyzopertha dominica*, *Sitophilus oryzae*, *Tribolium castaneum*, oviposition

Introduction

Cereals and pulses have great biological and nutritional value in developing countries, are lost upto 20-60 per cent by storage insect pests during storage (Arthur and Throne 2003; Babu et al. 2003; Shaaya et al. 1997). Post harvest deterioration causes economic losses due to obvious decay and adverse changes in the odour, taste, appearance and nutrition value (Phillips and Burkholder 1984; Mondal and Port 1994; Arlian et al. 1996). In addition, the arthropods transfer bacteria and microscopic fungi of pathogen importance on stick on their bodies or disseminate them via faeces (Wilbur and Mills 1978, Hubert et al. 2004). During recent years considerable attention has been paid towards exploitation of plant materials in protection of food commodities from insect infestations. Extracts of some plant species viz. *Lantana camara* (Saxena et al. 1992), *Illicium verum* (Ho et al. 1995), *Tithonia diversifolia* (Adedire and Akinneye, 2004) have been reported to possess strong insecticidal activity against different storage insects. Plant derived products namely, azadirachtin from *Azadirachta indica*, pyrethrin from *Chrysanthemum cinerariaefolium*, carvone from *Carum carvi* and allyl isothiocyanate from mustard and horseradish oil have been received global attention due to their pesticidal properties and potential to protect several food commodities (Hartmans et al. 1995; Ward 1998; Varma and Dubey 1999; Athanassiou et al. 2005). Essential oils produced by different plant genera have been reported to be biologically active and are endowed with insecticidal, antimicrobial and bio regulatory properties (Mishra and Dubey 1994; Varma and Dubey 1999; Dubey et al. 2004; Holley and Patel 2005). The volatility and biodegradability of flavour compounds of angiosperm will be advantageous if they are developed as pesticide insecticide (French 1985). There may be least chance of residual toxicity by treatment of food commodities with volatile substances of higher plant origin.

Aegle marmelos (L.) Correa (Rutaceae), commonly known as Bael, is a sacred tree for Hindu Religion, native to northern India, but is found widely throughout the

*Corresponding author. mailing address: Centre of Advanced Study in Botany, Faculty of Science, Banaras Hindu University, Varanasi- 221005, India. Phone: 91-542-2307146, 2307147, Fax: 91-542-268174, E- mail: nkubeybhu@gmail.com.

Indian peninsula and in Ceylon, Burma, Thailand and Indo-China (Bailey 1963). All parts of the tree namely,

root, leaf, trunk, fruit and seed are used for treatment of many different diseases. The constituents of *Aegle* are used in heart diseases (Kakiuchi et al. 1991), inflammatory and wound healing (Udupa et al. 1994). Leaves of *A. marmelos* have been reported as hypoglycemic effect (Santhoshkumari and Devi 1990; Sharma et al. 1996). The essential oil from the leaves of *A. marmelos* is known to exhibited antifungal properties (Renu et al. 1986; Rana et al. 1997).

Keeping these facts in mind, in the present investigation, essential oil of *A. marmelos* was tested as an *in vivo* fumigant to protect stored gram from insect pest *Callosobruchus chinensis* (L.) (Bruchidae) and wheat from *Rhyzopertha dominica* (F.), *Sitophilus oryzae* (L.) and *Tribolium castaneum* (Herbst). The effect of *Aegle* oil applications on mortality, oviposition and adult development of *C. chinensis* has also been determined.

Materials and Methods

Rearing of test insects. Rearing of *C. chinensis* was maintained on cowpea seeds (*Vigna unguiculata* (L.) Walp.) (Babu et al. 2003; Jenkins et al. 2003). *R. dominica*, and *S. oryzae* were maintained on wheat grains (*Triticum aestivum* (L.) while *T. castaneum* on wheat flour at $27 \pm 2^\circ\text{C}$ and RH $80 \pm 5\%$ (Perez-Mendoza et al., 2004). Forty adult insects were released separately in 200 g of commodities (wheat/cowpea/wheat flour) in plastic containers covered by muslin cloth. After 24 hours, adult insects were removed and the commodities were incubated in a temperature/humidity controlled cabinet ($27 \pm 2^\circ\text{C}$ and RH $80 \pm 5\%$) in darkness to obtain same aged insects. Adult insects were 2-4 days old when used in the bioassays.

Extraction of essential oil. Essential oil was extracted from fresh leaves of *A. marmelos* (300 g) through hydrodistillation by Clevenger's apparatus for 4 h. The extracted oil (yield 0.3 % v/w) was dried over anhydrous sodium sulphate and the essential oil was stored in clean glass vials.

Chemical characterization. The essential oil of *A. marmelos* leaves was subjected to GC-MS analysis. GC-MS analysis was performed on Perkin Elmer Turbomass/Auto XL Instrument (Perkin Elmer, Inc. USA) at 70 eV with EQUITY-5 capillary column (60m X 0.32 mm X 0.25 μm). Helium was used as carrier gas @ 10 psi column head pressure. Injector and detector temperature were at 220°C and 280°C , respectively. Split ratio was 1:50. The oven temperature was programmed from 100 to 280°C @ $3^\circ\text{C}/\text{min}$ with initial and final holdup time 2 min. The compounds were identified by comparing their retention times, mass spectra with authentic standards and the fragmentation patterns of the MS with NIST and Wiley GC-MS computer library of essential oils.

Fumigation of wheat and gram samples by the essential oil of *A. marmelos*. The essential oil of *A. marmelos* was used to fumigate the wheat and gram samples separately by the method adapted by Shaaya et al. (1997) and Kumar et al. (2007). Five hundred g of wheat samples (var. Malviya 234, 10.93 % moisture) were kept separately in closed plastic containers (35 cm diameter x 16 cm). Care was taken to use uninfested freshly harvested wheat grains. Twenty five individuals of each insect species, i.e. *R. dominica*, *S. oryzae* and *T. castaneum* of mixed sex were introduced in the containers. Requisite amount of the oil of *A. marmelos* was introduced separately in the plastic containers by soaking in cotton swab so as to procure concentration of 500 ppm. The containers were made airtight. The freshly harvested uninfested gram samples (var. PUSA 362, 10.75 % moisture) were also similarly provided with 25 individuals of *C. chinensis* and fumigated with the oil of *A. marmelos* at 500 ppm. The wheat and gram inoculated with the test insects without oil treatment served as controls. After 24 months of storage at laboratory conditions in a temperature/humidity control cabinet ($27 \pm 2^\circ\text{C}$ and RH $80 \pm 5\%$) in darkness. The efficacy of *A. marmelos* oil on insect infestation was determined by calculating grain damage (%), weight loss (%) and feeding deterrence (%) of treated and control sets. The grain damage was determined by counting feeding injuries and emergence holes on the surface of the grains. The weight loss (%) of wheat and gram samples in the treated and control sets was calculated by fresh weight basis using the formula suggested by Parkin (1956).

$$\text{Weight loss (\%)} = \frac{\text{WI} - \text{W}}{\text{WI}} \times 100$$

where WI and W represents the weight of grains before and after the experiment, respectively
Feeding deterrence was calculated using the feeding deterrent index following Isman (1990):

$$\text{Feeding deterrent index (FDI) [\%]} = \frac{\text{C} - \text{T}}{\text{C} + \text{T}} \times 100$$

where C and T is the weight loss in the controls and in the fumigated sets, respectively.

Effect of essential oil on mortality, oviposition and adult development of bruchids on cow pea. The essential oil of *A. marmelos* was tested for its *in vivo* effects on insect mortality, oviposition and adult emergence on *C. chinensis* following Bright et al. (2001) and Kumar et al. (2007). A stock solution of the essential oil was prepared by dissolving 100 μl of *Aegle* oil in 1.9 ml of acetone. Twenty five seeds of cow pea (*Vigna unguiculata* (L.) Walp.) were filled in glass vials (6.3 X 2 cm diameter) and treated separately with different dose i.e. 100, 10, 1.0 and 0.1 μl of the oil. The seeds were then dressed by continuous shaking for five minutes for proper mixing of the oils on the seeds. For control sets the seeds were dressed in requisite amount of acetone in place of the oil. The treated samples were kept in temperature-humidity control cabinet ($27 \pm 2^\circ\text{C}$ and

RH 80± 5%). After 24 hours five bruchids of mixed Requisite control sets were kept for each treatment set. After 24 hours the mortality of insects was observed in each vial and all insects (live and dead) were removed. The number of eggs laid on seeds of treated and control seeds were counted after three days of starting the experiment. The number of adult insects emerged in cow pea samples in each treated and control set was counted after 30, 45 and 90 days of storage.

Statistical analysis. All the experiments were repeated thrice and data was reported as mean value ± SE. The statistical analysis was performed by one way analysis of variance and means were compared by least significance difference test ($P < 0.05$) using the SPSS statistical software package (SPSS, ver. 10.0; Chicago, IL, USA). Further, the data was subjected to Student's 't' test to analyze the effect of *Aegle* oil on grain damage as well as weight loss of wheat and gram with control. The correlation coefficient was calculated between dose-mortality, dose-oviposition, dose-adult emergence, mortality-oviposition and oviposition-adult emergence using software Origin (Origin 6.0 Northampton, MA, USA).

Results

The GC-MS analysis of the *A. marmelos* oil depicted the presence of following compounds viz. α pinene (0.28 %), sabinene (0.14 %), limonene (88.57 %), ocimene (2.29 %) and p caryophyllene (0.06 %). Limonene was found to be major component in *Aegle* oil. As it evident from Table 1 that oil significantly protected stored gram from *C. chinensis* and wheat samples from *R. dominica* and *S. oryzae* for the first two years ($P < 0.05$; LSD). The feeding deterrent index of the oil for *C. chinensis*, *R. dominica* and *S. oryzae* was 91.51, 97.26 and 98.02 % respectively while it was -6.18 % against *T. castaneum*. There was 100 % grain damage in *T. castaneum* while 7.0, 3.67 and 1.67 % grain damage was found in *C. chinensis*. *R. dominica* and *S. oryzae* infested grains respectively. However, significant reduction in weight loss was found in fumigated gram and wheat against the test insects except *T. castaneum*.

The effect of *Aegle* oil on mortality, oviposition and adult development of *C. chinensis* in treated cowpea samples is presented in Table 2. *C. chinensis* showed 71.41 % mortality at 100 μ l dose of oil. Oviposition deterrent activity of the oil for *C. chinensis* enhanced with dose. The oviposition was reduced to 56.25 % at 100 μ l oil dose. The reduction of hatching was also directly proportional to oil dose. *Aegle* oil checked more than 70 % of adult emergence of *C. chinensis* at different doses.

There was positive relationship between dose of the oil and adult mortality of *C. chinensis* ($R = 0.898$; $P < 0.001$) i.e. mortality of bruchids increased with

sex were introduced in each vial separately, increasing dose of *Aegle* oil (Fig. 1a). The negative relationship was found between dose x oviposition ($R = -0.958$; $P < 0.0001$) and dose x adult emergence ($R = -0.6419$; $P < 0.05$) of *C. chinensis*, indicating enhancement of insect mortality and reduction in oviposition and adult emergence with increasing dose of *Aegle* oil (Fig. 1 b, c). Linear regression analysis showed the significant relationship between mortality and oviposition; and the number of eggs laid on cowpea and the corresponding number of emerging insects. Oviposition was directly proportional to adult emergence of bruchids while it was inversely proportional to mortality of the insects (Fig. 1 d, e).

Discussion

In the present study the essential oil of *A. marmelos* exhibited as botanical fumigant in protection of stored gram and wheat by enhancing feeding deterrence and reducing grain damage as well as weight loss of *C. chinensis*, *R. dominica* and *S. oryzae*. The findings are in accordance with Kumar et al. (2007) and Varma and Dubey (2001) who investigated that essential oil of *Cymbopogon martinii*, *Caesulia axillaris* and *Mentha arvensis* protected stored gram and wheat from *C. chinensis*, *S. oryzae* and *T. castaneum* for first 12 months of storage. In the present investigation the shelf life of the *Aegle* oil in protection of insect infestation was 24 months thus more than the oils reported earlier. Plant extracts and essential oils are known to possess repellent, ovicidal and insecticidal activities against various stored grain insects (Hill and Schoonhoven 1981; Desmarchelier 1994). The adult mortality might be attributed to the contact toxicity or to the abrasive effect on the pest cuticle (Mathur et al. 1985), which might also interfere with the respiratory mechanism of insect (Schoonhoven 1978; Agarwal et al. 1988; Kim et al. 2003). Results of present and earlier studies indicate that essential oil might be useful for managing coleopterous insects in enclosed spaces such as storage bins, glasshouse and buildings etc. because of their fumigant action. In addition, the interesting finding of the present study was the dissimilar performance of the *A. marmelos* oil against the test insects as well as food commodities. The same dose level of *A. marmelos* oil did not provide same level of protection against different commodities. The structure of the grains, texture, dust deposition and degree of adherence to the kernel affect the efficacy of insecticidal products (Subramanyam and Roseli 2000, Korunic 1997; Vayias and Athanassiou 2004). Thus, the dose should be prescribed according to the target species and the commodities.

In the present study the mortality, oviposition and adult emergence of *C. chinensis* were found vary significantly with different doses of *Aegle* oil.

Callosobruchus females generally prefer smooth seed varieties for oviposition (Haines 1991). Therefore, the cow pea was selected for oviposition and adult

Table1. Fumigant efficacy of *A. marmelos* oil on stored gram and wheat infested with four insect pests at 500 ppm

Essential oil	Gram			Wheat								
	<i>C. chinensis</i>			<i>R. dominica</i>			<i>S. oryzae</i>			<i>T. castaneum</i>		
	Grain Damage (%)	Weight loss (%)	FDI Index (%)	Grain Damage (%)	Weight loss (%)	FDI Index (%)	Grain Damage (%)	Weight loss (%)	FDI Index (%)	Grain Damage (%)	Weight loss (%)	FDI Index (%)
<i>A. marmelos</i>	7.00 ^a ± 0.30	2.64 ^a ± 0.21	91.51 ± 0.69	3.67 ^a ± 0.33	0.72 ^a ± 0.17	97.26 ± 0.66	1.67 ^a ± 0.30	0.49 ^a ± 0.09	98.02 ± 0.36	100.00 ^a ± 0.00	54.80 ^a ± 0.12	-6.18 ± 0.27
Control	100.00 ^b ± 0.00	59.50 ^b ± 0.29		100.00 ^b ± 0.00	52.07 ^b ± 0.46		100.00 ^b ± 0.00	48.96 ^b ± 0.15		100.00 ^a ± 0.00	48.45 ^b ± 0.32	

±: Standard error

Means within each column followed by the different letter are significantly different (P<0.05, Student's *t* test)

Table 2. Effect of *A. marmelos* oil on adult mortality, oviposition and adult emergence in *C. chinensis* on treated cow pea

Dose of essential oil (μ l)	Per cent adult mortality	Per cent reduction in egg number	Per cent reduction in adult Emergence (Days after treatment)		
			30 DAT	45 DAT	90 DAT
100	71.41 \pm 7.10 ^a	56.25 \pm 2.08 ^a	72.42 \pm 0.15 ^b	62.68 \pm 1.50 ^a	63.29 \pm 1.30 ^a
10	49.99 \pm 7.10 ^{ab}	47.92 \pm 2.40 ^a	70.90 \pm 1.82 ^{ab}	46.26 \pm 2.60 ^b	41.76 \pm 3.35 ^b
1.0	42.85 \pm 7.10 ^b	34.03 \pm 1.40 ^b	69.09 \pm 3.35 ^{ab}	31.30 \pm 3.00 ^c	37.97 \pm 3.34 ^b
0.1	14.28 \pm 0.00 ^c	22.22 \pm 3.02 ^c	65.45 \pm 1.82 ^a	25.36 \pm 1.50 ^c	25.31 \pm 3.30 ^c

\pm : Standard error

DAT: Days after treatment

Means within each column followed by the different letter are significantly different ($P < 0.05$, LSD).




FIGURE. 1a. Effect of dose of *Aegle* oil on adult mortality of *C. chinensis*




FIGURE 1C. Effect of dose of *Aegle* oil on adult development of *C. chinensis*




FIGURE 1d. Relationship between adult mortality and oviposition of *C. chinensis*




FIGURE. 1e. Relationship between number of eggs of *C. chinensis* laid on cowpea and number of insects hatched

emergence trails in place of gram. *C. chinensis* females laid significantly fewer eggs on cowpea treated with different doses of *Aegle* oil than control sets, indicating that oil deterred *C. chinensis* females from ovipositing. Furthermore fewer insects were hatched in treated cowpea than the control at 30, 45 and 90 DAT (days after treatment). Since *C. chinensis* females deposit their eggs on the seed surface (Haines 1991), neonate larvae have to penetrate through the outer skin of the cowpea to reach the endosperm where the biologically active compounds from the *Aegle* oil might have deterred them. The reduction in adult emergence could either be due to egg-mortality or larval mortality or even reduction in hatching of the eggs. Oviposition inhibitors have the advantage of attacking a pest at the start of its life cycle. The insect is deterred from laying its eggs on the foodstuff, thus preventing the pest population from increasing.

The insecticidal constituents of many essential oils are mainly monoterpenoids (Coats et al. 1991; Konstantopoulou et al. 1992; Regnault-Roger and Hamraoui 1995; Kim et al. 2003). Monoterpenoids have been reported to inhibit reproduction of stored insects at several steps of the cycle: inhibition in oviposition, ovicidal effects, a larvicidal effect on neonate larvae before the penetration of the seeds or a larvicidal effect on larvae settled within the seed, thus inhibiting the emergence of imagoes. A strong

inhibition of oviposition was produced by carvacrol, thymol and terpineol. α pinene strongly inhibited the larval penetration and less effective to oviposition inhibition against the storage insect pest, *Acanthoscelides obtectus* (Regnault-Roger and Hamraoui, 1995). In the present study limonene was the major component of the oil of *A. marmelos* and it may be responsible for fumigant toxicity, oviposition deterrent, inhibition of adult development, and feeding-deterrent activities of test insects. The mode of toxicity for monoterpenoids is believed to be via competitive inhibition of acetylcholinesterase (Ryan and Byrne 1988).

Compounds with feeding deterrents are generally toxic to insects or cause physiological disturbances of development or oviposition (Nawrot and Harmantha 1994). For example azadirachtin is well known antifeedant and also interferes with insect growth (Siddig 1980). *A. marmelos* oil was mixture of a number of components that would reduce the chances for the development of insect resistance, which could be due to the diffusion of the gene selection process (Begon et al. 1999; Davies 1992). Although animal toxicity with *A. marmelos* oil were not conducted in the present study, however, concerns for residue of essential oil pesticides on food grains should be mitigated by the growing body of evidence that many essential constituents acquired through the diet are

actually beneficial to human health (Huang et al. 1994).

In summary, *Aegle* oil may be used as botanical insecticide against different stored grain insect pests causing infestation in stored wheat and pulses. The *Aegle* oil enhanced feeding deterrence of *C. chinensis*, *R. dominica* and *S. oryzae*. Therefore, insects were incapable to infest grain and to cause grain damage. *A. marmelos* significantly reduced oviposition and adult emergence of *C. chinensis* in treated cowpea. Moreover, because of the use in traditional medicine in cure of different human diseases, the *Aegle* oil may be used as semiochemicals mediating phytopesticide to protect stored food commodities in developing countries, for which some farmers may not have easy access to chemical insecticides.

Acknowledgements

Authors are thankful to CSIR, New Delhi for financial assistance and head, Department of Botany, Banaras Hindu University for providing laboratory facilities.

References

- Adedire CO, Akinneye JO. 2004. Biological activity of tree marigold, *Tithonia diversifolia*, on cowpea seed bruchid, *Callosobruchus maculatus* (Coleoptera: Bruchidae). *Ann. Appl. Biol.* 144:185-189.
- Agarwal A, Lal S, Gupta KC. 1988. Natural products as protectants of pulse beetles. *Bull. Grain Tech.* 26:154-164.
- Arthur FH, Throne JE. 2003. Efficacy of diatomaceous earth to control internal infestations of rice weevil and maize weevil (Coleoptera: Curculionidae). *J. Econ. Entomol.* 96:510-518.
- Arlian, LG. 2002. Arthropod allergens and human health. *Annu. Rev. Entomol.* 47:395-433.
- Athanassiou CG, Kavallieratos NG. 2005. Insecticidal effect and adherence of PyriSec® in different grain commodities. *Crop Prot.* 24:703-710.
- Athanassiou CG, Kontodimas DK, Kavallieratos NG, Anagnou-Veroniki M. 2005. Insecticidal effect of NeemAzal against three stored product beetle species on rye and oats. *J. Econ. Entomol.* 98:1733-1738.
- Babu A, Hern A, Dorn S. 2003. Sources of semiochemicals mediating host findings in *Callosobruchus chinensis* (Coleoptera: Bruchidae). *Bull. Entomol. Res.* 93:187-192.
- Bailey LH. 1963. *How Plants Get Their Names*. Dover Publications, Inc., New York, New York.
- Begon M, Harper JL, Townsend CR. 1999. *Ecology: individuals, populations and communities*. Blackwell Scientific Publications, Oxford, UK., pp 551-582.
- Bright AA, Babu A, Ignacimuthu A, Dorn S. 2001. Efficacy of crude extracts of *Andrographis paniculata* Nees. on *Callosobruchus chinensis* L. during post harvest of cowpea. *Int. J. Exp. Biol.* 39:715-718.
- Coats JR, Karr LL, Drewes CD. 1991. Toxicity and neurotoxic effects of monoterpenoids in insects and earthworms. In: *Naturally Occurring Pest Bioregulators*. Hedin PA (ed.) ACS Symposium Series No. 449, American Chemical Society, Washington, DC., pp 305-316.
- Davies WP. 1992. Prospects for pest resistance to pesticides, *Pest Management and the Environment in 2000*. In: Kadir AASA, Barlow HS (Eds), CAB International, Wallingford, UK., pp 95-110.
- Desmarchelier JM. 1994. Grain protectants: trends and developments. In: *Stored Product Protection Vol. 2*. Highley E, Wright EJ, Banks HJ, Champ BR (Eds) CAB International, Wallingford, UK. pp 722-728.
- Dubey NK, Kumar R, Tipathi P. 2004. Global Promotion of herbal medicine: India's opportunity. *Curr. Sci. (India)* 86:37-41.
- French RC. 1985. The bioregulatory action of flavour compounds on fungal spores and other propagules. *Annu. Rev. Phytopathol.* 23:173-199.
- Haines CP. 1991. *Insects and arachnids of tropical stored products: their biology and identification. A training manual*, 2nd ed. Natural Resources Institute. Hobbs, Southampton, UK.
- Hartmans KJ, Diepenhorst P, Bakker W, Gorris LGM. 1995. The use of carvone in agriculture, sprout suppression of potatoes and antifungal activity against potato tuber and other plant diseases. *Ind. Crop Prod.* 4:3-13.
- Hill JM, Schoonhoven AV, 1981. The use of vegetable oils in controlling insect infestations in stored grains and pulses. *Recent Advan. Food Sci. Technol.* 1: 473-481.
- Ho SH, Ma Y, Goh PM, Sim KY. 1995. Star anise, *Illicium verum* Hook f. as a potential grain protectant against *Tribolium castaneum* Herbst and *Sitophilus zeamais* Motsch. *Postharvest. Biol. Tech.* 6:341-347.
- Holley AH, Patel H. 2005. Improvement in shelf life and safety of perishable foods by plant essential oils and smoke antimicrobials. *Int. J. Food Microbiol.* 22:273-292.
- Huang MT, Ferraro T, Ho CT. 1994. Cancer chemoprevention by phytochemicals in fruits and vegetables. *American Chemical Society Symposium Series* 546: 2-15.
- Hubert J, Stejskal V, Munzbergova Z, Kubatova A, Vanova M, Zd'arkova E. 2004. Mites and fungi in heavily infested stores in the Czech Republic. *J. Econ. Entomol.* 97(6):2144-53.

- Isman MB, Koul O, Luezyński N. 1990. Insecticidal and antifeedant bioactivity of neem oil and their relationship to azadirachtin content. *J. Agric. Food Chem.* 38:1406-1411.
- Jenkins DA, Dunkel FV, Gamby KT. 2003. Storage temperature of Neem kernel extract: differential effects on oviposition deterrence and larval toxicity of *Callosobruchus maeulatus* (F.) (Coleoptera: Bruchidae). *Environ. Entomol.* 32:1283-1289.
- Kakiuchi N, Senaratne LRE, Huang SL, Yang XW, Hattori M, Pilapitiya U, Namba T. 1991. Effects of constituents of Beli (*Aegle marmelos*) on spontaneous beating and calcium-paradox of myocardial cells. *Plant Med.* 57: 43-46.
- Kavallieratos NG, Athanassiou CG, Pashalidou FG, Andris NS, Tomanović, Ž. 2005. Influence of grain type on the insecticidal efficacy of two diatomaceous earth formulations against *Rhyzopertha dominica* (F) (Coleoptera: Bostrychidae). *Pest Manag. Sci.* 61: 660-666.
- Kim SI, Roh JY, Kim DH, Lee HS, Ahn YJ. 2003. Insecticidal activities of aromatic plant extracts and essential oils against *Sitophilus oryzae* and *Callosobruchus chinensis*. *J. Stored Prod. Res.* 39: 293-303.
- Konstantopoulou L, Vassilopoulou L, Mauragani-Tsipidov P, Scouras ZG. 1992. Insecticidal effects of essential oils. A study of the effects of essential oils extracted from eleven Greek aromatic plants on *D. auraria*. *Experientia* 48:616-619.
- Korunic Z. 1997. Rapid assessment of the insecticidal value of diatomaceous earths without conducting bioassays. *J. Stored Prod. Res.* 33: 219-229.
- Kumar R, Srivastava M, Dubey NK. 2007. Evaluation of *Cymbopogon martinii* oil extract for control of post harvest insect deterioration in cereal and pulse. *J. Food Prot.* 70: 172-178.
- Mathur YK, Shankar K, Ram S. 1985. Evaluation of some grain protectants against *Callosobruchus chinensis* (L.) on black gram. *Bull. Grain Tech.* 23: 253-259.
- Mishra AK, Dubey NK 1994. Evaluation of some essential oils for their toxicity against fungi causing deterioration of stored food commodities. *Appl. Environ. Microbiol.* 60: 1401-1405.
- Mondal KAMSH, Port GR. 1994. Pheromones of *Tribolium* spp and their potential in pest management. *Agricul. Zool. Rev.* 6:121-148.
- Nawrot J, Harmatha J. 1994. Natural products as antifeedants against stored products insects. *Postharvest News and Information* 5:17N-21 N.
- Parkin EA. 1956. Stored Product Entomology (the assessment and reduction of losses caused by insects to stored food stuffs). *Annu. Rev. Entomol.* 1:233-240.
- Perez-Mendoza J, Throne JE, Dowell FE, Baker JE. 2004. Chronological age-grading of three species of stored-product beetles by using near-infrared spectroscopy. *J. Econ. Entomol.* 97:1159-1167.
- Phillips JK, Burkholder WE. 1984. Health hazards of insects and mites in food. *Insect Management for Food Storage and Processing*, In: Braur F (ed.) *The American Association of Cereal Chemists*, pp. 280-292.
- Rana BK, Singh UP, Taneja V. 1997. Antifungal activity and kinetics of inhibition by essential oil isolated from leaves of *Aegle marmelos*. *J. Ethnopharmacol* 57: 29-34.
- Renu HVM, Dubey NK, Dixit SN. 1986. Mycotoxic properties of the essential oil of *Aegle marmelos*. *Beit. Biol. Pflanz* 60:325-332.
- Regnault-Roger C, Hamraoui A. 1995. Fumigant toxic activity and reproductive inhibition induced by monoterpenes on *Acanthoscelides obtectus* (Say) (Coleoptera), a bruchid of kidney bean (*Phaseolus vulgaris* L.). *J. Stored Prod. Res.* 31:291-299.
- Ryan MF, Byrne O. 1988. Plant insect coevolution and inhibition of acetylcholinesterase. *J. Chem. Ecol.* 14:1965-1975.
- Santhoshkumari KS, Devi KS. 1990. Hypoglycemic effect of a new medicinal plants. *Ancient Science of Life* IX/4:221-223.
- Saxena RC, Dixit OP, Harshan V. 1992. Insecticidal action of *Lantana camara* against *Callosobruchus chinensis* (Coleoptera: Bruchidae). *J Stored Prod. Res.* 28:279-281.
- Schoonhoven AV. 1978. The use of vegetable oils to protect stored beans from bruchid attack. *J. Econ. Entomol.* 71:254-256.
- Shaaya E, Kostjukovski M, Eilberg I, Sukprakarn C. 1997. Plant oils as fumigants and contact insecticides for the control of stored-product insects. *J. Stored Prod. Res.* 33: 7-15.
- Sharma SR, Dwivedi SK, Varshney VP, Swarup D. 1996. Antihypoglycemic and Insulin release effects of *Aegle marmelos* leaves in streptozotocin diabetic rats. *Phytother. Res.* 10: 426-428.
- Siddig SA. 1980. Efficacy and persistence of powered neem seeds for treatment of stored wheat against *Trogogramma granarium*. In: Schmitterer H, Ascher KRS, Rembold, H (eds.) *Proceedings of the 1st International Neem Conference, Rottach-Egern, FRG.*, pp 251-257.
- Subramanyam Bh, Roesli R. 2000. Inert dusts. In: Subramanyam Bh, Hagstrum DW (Eds) *Alternatives to pesticides in stored-product IPM*. Kluwer Academic Publishers, Dordrecht., pp 321-380.
- Udupa SL, Udupa AL, Kulkarni DR. 1994. Studies on the anti-inflammatory and wound healing properties of *Moringa oleifera* and *Aegle marmelos*. *Fitoterapia* 65:119-123.

- Varma J, Dubey NK. 2001. Efficacy of essential oils of *Caesulia axillaris* and *Mentha arvensis* against some storage pests causing biodeterioration of food commodities. *Int. J. Food Microbiol.* 68:207-210.
- Varma J, Dubey NK. 1999. Prospectives of botanical and microbial products as pesticides of tomorrow. *Curr Sci (India)* 76:172-179.
- Vayias BJ, Athanassiou CG. 2004. Factors affecting efficacy of the diatomaceous earth formulation SilicoSec against adults and larvae of the confused beetle *Tribolium confusum* du Val (Coleoptera: Tenebrionidae). *Crop Prot.* 23:565–573.
- Ward SM, Delaquis PJ, Holley RA, Mazza G. 1998. Inhibition of spoilage and pathogenic bacteria on agar and pre-cooked roasted beef by volatile horse radish distillates. *Food Res. Int.* 31:19–26.
- Wilbur DA, Mills RB. 1978. Stored Grain Insects. In: Pfadt RE (Ed), *Fundamentals of Applied Entomology*, Macmillan Publishing Company, New York USA.

